

Orbiting Light

Orbiting Light

john.erich.ebner@gmail.com
<http://blackholeformulas.com>

September 14, 2014

Abstract

This paper postulates that light will orbit any great enough mass. We know that gravity deflects light from numerous cases of gravitational lensing so it is not too big a step to see that a deflection of light with one mass could become an orbit of light with the much greater mass of the Cosmos. We explore this simple classical model of energy and light in orbit with the black hole and the Cosmos in dynamic equilibrium. Our purpose is to solve the problems of the spiral shape and flat rotation curves of galaxies, prevalence of dark matter in galaxy dynamics and the source, uniformity and Planck satellite anomaly of the cosmic microwave background. Wiki calls these "unsolved problems of physics." They are shown to be a consequence of this model of the dynamic universe. The age, radius, mass, expansion rate and density of the universe also come from this model and are unambiguously shown to apply to a black hole and are in the range or magnitude of mainstream values frequently quoted. There are no infinities, singularities or free parameters which might be adjusted to reflect a point of view.

Key Words

Orbiting light or energy, expanding and rotating universe, gravity, dark matter, cosmic microwave background, unsolved problems in physics

Authors Note

This document was written with Latex <http://latex-project.org/ftp.html> and TexStudio <http://texstudio.sourceforge.net/>, both of which are excellent, open-source and free. The PDF pages it produces can be read in two page view and printed two pages at a time in landscape to save paper or make a book.

Contents

1	Hubble	6
2	Light in Orbit	8
3	Geons	9
4	Expanding Black Holes	10
5	The Cosmos rotates and expands	12
6	Variable G	13
7	Orbital forces decrease with time	14
8	Moving clocks or clocks in a gravitational field	15
9	Black Hole Density	16
10	Density compared to a Neutron Star using (9.4)	17
11	Merging Black Holes	18
12	Large Scale Structure	19
13	Spherical caps of merging spheres	19
14	Comparing Traditional Black Holes and Orbiting Light Black Holes	20
15	Tests for Orbiting Light Black Holes	23
16	Rotation of the Cosmos	23
17	Cosmic Expansion	28

18 Galactic Matters	30
19 Flat Rotation Curves of Galaxies	32
20 Galactic Expansion	33
21 Dark matter	34
22 Hubble expansion in the Solar System	35
23 Ellipses	39
24 Cosmic Cross Section	39
25 Blackbody watts from temperature	41
26 Uniformity of the CMB	42
27 Black body watts from the CMB	43
28 What is reasonable?	46

List of Figures

1	Expanding Cosmos	12
2	Triangular definitions	12
3	Merging black holes	18
4	Binary rotations	24
5	Logarithmic spirals	29
6	Galaxy expansion	33
7	Solar system expansion	36
8	Cosmic cross section	40
9	Expanding ellipses	41

1 Hubble

(age, radius, radial velocity)

Edwin Hubble[1] determined from the linear Doppler redshift of galaxies that they are receding at a rate proportional to distance. It is called Doppler because of the familiar frequency shift of sound with the velocity of the sound source. Here the frequency shift is in the light toward the red with increasing velocity of recession. The velocity of recession at a certain distance divided by that distance is a constant. In our examples:

$$\text{Hubble's constant} = \frac{\text{velocity}}{\text{distance}} = H_0 = 65 \frac{\text{km}}{\text{Mpc}} =$$

Kilometers per second per million parsecs, in smaller units this is:

$$H_0 = \frac{m}{s} \text{ per meter} = 2.11E-18 \frac{1}{s} \quad (1.1)$$

Using these smaller more familiar units may lead you to wonder if the expansion is also local. Does the Hubble expansion extend to the galaxy, to the solar system and to atoms? If the local expansion exists, was it missed because the size of the expansion is so small? Hubble's constant has units of 1/seconds which is frequency, angular velocity or 1/age. The reciprocal of Hubble's constant is:

$$\frac{1}{H_0} = 4.74E17 \text{ s} = 15 \text{ billion years} = \text{age} \quad (1.2)$$

We will use 15 billion years as the age in our examples hereafter called the age of the Cosmos[2]. Hubble's constant can be written:

$$H_0 = \frac{1}{\text{age}} = \frac{c}{c * \text{age}} = \frac{c}{15 \text{ billion light years}} \quad (1.3)$$

This implies that the Cosmos is expanding at the speed of light at its current perimeter,

$$r = c * \text{age} = 15 \text{ billion light years} = 1.42E26 \text{ m} \quad (1.4)$$

the current radius of the Cosmos[3]. This paper does not require or support space-time, the expansion of the vacuum of space or inflation so the radius is the radius. Hubble varies, but the radial velocity of expansion is constant, with distance divided by age at the perimeter.

$$radial\ velocity = vr = \frac{r}{age} = \frac{c * age}{age} = c \quad (1.5)$$

This is age, radius and radial velocity based on the expansion rate, not an affirmation of a creation event 15 billion years ago.

Vary Hubble's Constant

$$130 \frac{km}{s * Mpc} = \frac{1}{7.5\ billion\ years} = H_0 * 2 \quad (1.6)$$

$$70.4 \frac{km}{s * Mpc} = \frac{1}{13.89\ billion\ years} = H_0\ per\ WMAP\ satellite \quad (1.7)$$

$$67.15 \frac{km}{s * Mpc} = \frac{1}{14.56\ billion\ years} = H_0\ per\ Planck\ satellite \quad (1.8)$$

$$65 \frac{km}{s * Mpc} = \frac{1}{15\ billion\ years} = H_0\ per\ our\ examples \quad (1.9)$$

$$32.5 \frac{km}{s * Mpc} = \frac{1}{30\ billion\ years} = \frac{H_0}{2} \quad (1.10)$$

$$65 \frac{km}{s * Mpc} = \frac{1}{60\ billion\ years} = \frac{H_0}{4} \quad (1.11)$$

This suggests that at earlier times, Hubble's constant was bigger and the Cosmos was smaller. It is likely that some part

of the Hubble constant is due to non-doppler redshift [4]. If half of Hubble's constant is due to non-Doppler redshift then the Doppler portion is $32.5 \frac{km}{s * Mpc}$ and the age is 30 billion years not 15 billion years. The calculated age, mass and radius of the Cosmos increases with the decrease in the Doppler portion of the redshift. Therefore, if all the redshift is non-Doppler then the Cosmos is infinitely old and massive.

2 Light in Orbit

$$\frac{m * vt^2}{r} = \frac{G * m * M}{r^2} \quad (2.1)$$

In orbiting systems with two masses the centrifugal force equals the gravitational force. If the centrifugal force is stronger the bodies drift apart. If the gravitational force is stronger the bodies drift together. The average forces must be equal for the orbits to endure. vt is the tangent orbital velocity. M and m are masses with the radius r between them. G is the gravitational constant.

$$\frac{m * c^2}{r} = \frac{G * m * M}{r^2} \quad (2.2)$$

Substituted the tangent velocity $vt = c$. Now the orbital velocity is c and light can orbit at the radius of the Cosmos at $r = c * \text{age}$. We know that gravity deflects light from numerous cases of gravitational lensing. It is almost trivial to postulate that a deflection of light with one mass could become an orbit of light with a much greater mass.

$$m * c^2 = \frac{G * m * M}{r} \quad (2.3)$$

Multiplied by r . The orbiting rest energy equals the gravitational energy. The orbital velocity is c where light orbits at the specific radius r . Light might follow a circular orbital path, a bent path as in gravitational lensing or a spiral path with a decreasing gravitational force. Light could form a spherical shell. Any

mass or energy inside the shell would contribute to the total mass increasing the radius of the shell. Light could form an empty shell with energy confined to the shell since light has energy and consequently mass. An empty shell of a black hole with only orbiting light does requires some thought. Multiply by r/m . These three equations (2.4 - 2.6) are our definitions of a black hole.

$$c^2 * r = G * M \quad (2.4)$$

$$\frac{c^2 * r}{G * M} = 1 \quad (2.5)$$

$$\frac{M}{r} = \frac{c^2}{G} = 1.35E27 \frac{kg}{m} = \frac{mass}{radius} \text{ ratio} \quad (2.6)$$

The current value of the mass per radius ratio. Black hole mass increases by $1.35E27 \frac{kg}{meter}$. This is a solar mass every 1476 meters.

3 Geons

In 1955, John Archibald Wheeler found an interesting way to treat the concept of body in general relativity which he called Geons[5].

”An object can, in principle, be constructed out of gravitational radiation or electromagnetic radiation, or a mixture of the two, and may hold itself together by its own gravitational attraction. The gravitational acceleration needed to hold the radiation in a circular orbit of radius r is of the order of $\frac{c^2}{r}$. The acceleration available from the gravitational pull of a concentration of radiant energy of mass M is of the order $\frac{GM}{r^2}$. The two accelerations agree in order of magnitude when ...”

$$\frac{c^2}{r} = \frac{GM}{r^2} \text{ or } c^2 * r = G * M \quad (3.1)$$

"... A collection of radiation held together in this way is called a geon (gravitational electromagnetic entity) and is a purely a classical object. ... Studied from a distance, such an object presents the same kind of gravitational attraction as any other mass... A geon made of pure gravitational radiation a ... gravitational geon ... owes its existence to a localized – but everywhere regular – curvature of spacetime, and to nothing more. In brief, a geon is a collection of electromagnetic or gravitational wave energy, or a mixture of the two, held together by its own gravitational attraction, that describes mass without mass." [6]

Wheeler gave the name to the "black hole" in 1967. Curious that he did not use the formula he gave to geons in 1955 and apply it to black holes.

4 Expanding Black Holes

$$r = vr * age = c * age = \frac{M * G}{c^2} \text{ or } G = \frac{c^3 * age}{Mc} \quad (4.1)$$

If $r = vr * age$ varies with age, as it must in an expanding Cosmos, then $G = \frac{c^3 * age}{Mc}$ must also vary with age. M is the mass of any black hole. Mc is the mass of the Cosmos. Mass and c are constants. vr is the constant radial velocity of expansion at its perimeter.

$$vr * age = \frac{M * c^3 * age}{Mc * c^2} \quad (4.2)$$

Substituted for $r = vr * age$ and $G = \frac{c^3 * age}{Mc}$. The age cancels.

$$vr = \frac{M * c}{Mc} \quad (4.3)$$

The radial velocity of expansion is constant and proportional to mass.

$$vr = M * 1.568E-45 \frac{m}{s * kg} \quad (4.4)$$

$$\frac{vr}{c} = \frac{M}{Mc} \quad (4.5)$$

The radial velocities and masses are ratios.

Black hole mass and radial velocity using (4.4)

$$Ten \text{ solar masses} = 1.99E31 \text{ kg so } vr = 3.12E-14 \frac{m}{s} \quad (4.6)$$

$$A \text{ billion solar masses} = 1.99E39 \text{ kg so } vr = 3.12E-6 \frac{m}{s} \text{ and}$$

$$a \text{ billion solar mass black hole expands at } 98 \frac{m}{year} \quad (4.7)$$

$$9.61E22 \text{ solar masses} = \text{the mass of the Cosmos} = Mc \text{ and} \quad (4.8)$$

$$= 1.91E53 \text{ kg so } vr = 299792458 \frac{m}{s} = c \quad (4.9)$$

Black holes expand in proportion to their mass. The radius of the Cosmos expands at the speed of light c , or 1 light year per year, because it has enough mass for it to expand at c . This proportionally small expansion is one year in the age of the Cosmos or one part in 15 billion per year.

Figure 1: Expanding Cosmos





Figure 2: Triangular definitions



5 The Cosmos rotates and expands

The Cosmos is expanding and rotating in figure [1]. See the animation at [9]. Each circle is a cross section of the Cosmos at a different time. Observers would look out in any direction and see the Cosmos expanding faster at greater distances. This is a Hubble expansion. Light orbits at the perimeter of the Cosmos. As the Cosmos increases in size, light orbits farther out, and the rotation slows down.

The Cosmos is rotating and blowing up like a balloon but very slowly in relation to its size as shown in figure [2]. See the animation at [10]. Like a spinning ice skater slows her spin by extending her arms, the spinning universe slows in its rotation while expanding. We rotate with respect to that which is very far away and is outside our dynamic unit Cosmos. The triangles show the way it expands and rotates. The length of the hypotenuse of the triangle is the velocity of light times the age of the Cosmos, at that particular time, or $r = c \cdot \text{age}$. The hypotenuse expands and has a radial and tangent velocity of the speed of light $v = c$. The base of the triangle expands and ro-

tates at its fixed fraction of the Cosmos. This fraction fr is the cosine of the triangle. The circled point on the base of the triangle is where we might be located. The location stays at the same constant fraction fr of the radius of the Cosmos $r = fr * c * age$ and constant velocity $v = fr * c$. Here fr at the location is $.7$ and the velocity is $v = fr * c = .7 * c$. This is a Hubble expanding Cosmos. Every location within this Cosmos has a constant tangent and radial velocity which are proportional to its radius. Since the velocity is constant, the acceleration is zero and no force and no power is required for it to continue on its journey. A location spirals out as the Cosmos expands and rotates.

6 Variable G

You may have noticed that G has a value of about $1/15$ billion.

$$G = 6.673E-11 \frac{m^3}{kg * s^2} \approx \frac{1}{15 \text{ billion}} \frac{m^3}{kg * s^2} \quad (6.1)$$

G increases by a small amount every year.

$$\frac{\Delta G}{year} = \frac{1}{15 \text{ billion}} - \frac{1}{15 \text{ billion} + 1} = 4.5E-21 \frac{m^3}{kg * s^2} \quad (6.2)$$

G increases by a small amount each second.

$$\frac{\Delta G}{second} = \frac{c^3 * one \ second}{Mc} = 1.41E-28 \frac{m^3}{kg * s^2} \quad (6.3)$$

$$\frac{\Delta G}{year} = \frac{c^3 * seconds \ in \ year}{Mc} = 4.44E-21 \frac{m^3}{kg * s^2} \quad (6.4)$$

$$\frac{\Delta G}{age} = G = \frac{c^3 * seconds \ in \ age}{Mc} = 6.673E-11 \frac{m^3}{kg * s^2} \quad (6.5)$$

How do we detect such a small ongoing increase in the gravitational constant?

$$c^2 * r = G * M \text{ Our black hole definition from (2.4)} \quad (6.6)$$

We take c and M to be constants. If r is a variable and increases with age then G is a variable and must increase with age. If G and r are both proportional to age then the orbiting and gravitational energy are constant as the black hole expands. Energy is conserved. This can be written $G = gk * \text{age}$. gk is the increase in the gravitational constant per second as seen above.

$$gk = \frac{G}{\text{age}} = G * H_0 = 1.41E-28 \frac{m^3}{kg * s^3} \quad (6.7)$$

gk appears to be a universal constant in contrast to H_0 which is Hubble's variable constant and G which is the variable gravitational constant.

$$c^2 * r = G * M \text{ Our black hole definition from (2.4)}$$

$$c^3 = gk * M \quad (6.8)$$

Substituted for $r = c * \text{age}$, $G = gk * \text{age}$. Multiply by 1/age.

$$Mc = \frac{c^3}{gk} = 1.91E53 \text{ kg The mass of the Cosmos [7]} \quad (6.9)$$

6E56 g or 6E53 kg according to Ciufolini and Wheeler[8]. If mass is due to gravity rather than gravity due to mass, would it explain rest energy equals gravitational energy in (2.3)? Gravity is likely due to charge. See Electric Gravity.[11]

7 Orbital forces decrease with time

The orbital centrifugal and gravitational forces are:

$$\frac{m * v^2}{r} = \frac{G * m * M}{r^2} \quad (7.1)$$

$$\frac{m * c^2}{c * age} = \frac{gk * age * m * c^2}{gk * (c * age)^2} \quad (7.2)$$

Substituted for $v = c$, $r = c * age$, $G = gk * age$ and for $M = Mc = c^3 / gk$.

$$\frac{m * c}{age} = \frac{m * c}{age} \quad (7.3)$$

The centrifugal and gravitational forces are equal and decrease with age. The forces were much stronger in the past. The stars burned brighter. Light from a distant earlier time has a greater redshift. How does this affect Hubble's constant and distance calculations? Star formation, accretion and fusion rates would have been much faster with the stronger forces in the past. The standard candle, Type 1A supernovas, would have been affected. What does this do to the accelerating universe dark energy theory?

8 Moving clocks or clocks in a gravitational field

run slower and atoms emit lower frequency "red shifted" radiation. Atoms expand or contract with changing kinetic energy [12]. Consider a Bohr atom with an orbiting electron and proton. When the Bohr atom is accelerated it acquires kinetic energy. The kinetic energy is stored in the atom as an increase in mass.

$$m \uparrow * v \downarrow = \text{constant momentum} \quad (8.1)$$

When the mass within the atom goes up, the orbital velocity within the atom goes down. Further, by analogy with the laws of Kepler.

$$\frac{m * v^2}{r} = \frac{G * m * M}{r^2} \text{ or } v^2 * r = G * M \quad (8.2)$$

$$v^2 \downarrow * r \uparrow = \text{constant} \quad (8.3)$$

When the orbital velocity within the atom goes down, the radius of the Bohr atom goes up. In quantum mechanics, we see that an expansion of the Bohr radius causes an increase in the wave length of the emitted "red shifted" radiation and a decrease in the atomic energy levels.

9 Black Hole Density

$$\text{mass of the Cosmos} = \frac{c^2 * r}{G} = \frac{c^2 * c * \text{age}}{gk * \text{age}} = \frac{c^3}{gk} \quad (9.1)$$

$$\text{radius of the Cosmos} = \frac{\text{mass} * G}{c^2} = \frac{c^3 * gk * \text{age}}{gk * c^2} = c * \text{age} \quad (9.2)$$

$$\text{surface area} = \frac{\text{mass}^2 * 4\pi * G^2}{c^4} = 4\pi * c^2 * \text{age}^2 \quad (9.3)$$

$$\text{density} = \frac{\text{mass}}{\text{volume}} = \frac{3 * \text{mass}}{4\pi * r^3} = \frac{3 * c^6}{4\pi * \text{mass}^2 * G}$$

Substitute for the mass of the Cosmos = $\frac{c^3}{gk}$

$$\frac{3}{4\pi * gk * \text{age}^3} = 1.6E-26 \frac{kg}{m^3} = 1.6E-29 \frac{g}{cm^3} \quad (9.4)$$

The lowest density black hole is our Cosmos. The mass of a proton or hydrogen atom is 1.67E-27 kg so the average density of the Cosmos [10] is about ten protons per cubic meter and is decreasing with the cube of the age of the Cosmos.

10 Density compared to a Neutron Star using (9.4)

$$20 \text{ solar mass black hole} = 3.68E17 \frac{kg}{m^3} \text{ and } r = 29531m \quad (10.1)$$

$$10 \text{ solar mass black hole} = 1.47E18 \frac{kg}{m^3} \text{ and } r = 14765m \quad (10.2)$$

$$5 \text{ solar mass black hole} = 5.9E18 \frac{kg}{m^3} \text{ and } r = 7383m \quad (10.3)$$

$$1.4 \text{ solar mass black hole} = 7.52E19 \frac{kg}{m^3} \text{ and } r = 2067m \quad (10.4)$$

$$1.4 \text{ solar mass neutron star} = 1E21 \frac{kg}{m^3} \text{ and } r = 874m \quad (10.5)$$

The density increases as the mass of the black hole decreases up to a limit and then the black hole is seen as a neutron star not a black hole. Less than five solar masses is observable as a neutron star not a black hole. These are in the range of the nuclear density [11] as calculated by particle physicist.

$$\text{Nuclear density} = 1 * 10E21 = 1E21 \frac{kg}{m^3} \quad (10.6)$$

Astronomers say neutron stars cluster around a neutron star mass [12] of 1.4 solar masses = 2.8E30 kg, and x-ray burster radius [13] of 9600 - 11000 m in x-ray bursters yields,

$$x - \text{ray burster density} \approx 5.0E17 \frac{kg}{m^3} \Rightarrow 7.6E17 \frac{kg}{m^3} \quad (10.7)$$

Figure 3: Merging black holes



We will use nuclear density $1E21 \frac{kg}{m^3}$ as the highest density in neutron stars. The highest density in black holes is therefore less than nuclear density in neutron stars. The density decreases as the cube of the radius or square of the mass so a larger radius or mass means a lower density. As the mass goes up the density goes down. We have black holes without infinities.

11 Merging Black Holes

Looking at figure [3]. Visualize two spheres that overlap, their contents merge, thereafter being enclosed by a single larger sphere, of their combined diameters. The light and energy in orbit around each black hole, will after merging, and after their travel through space, eventually orbit at this new larger radius. The light that previously orbited around each black hole will combine to orbit at the new combined black hole radius. All these photons orbiting at the same radius have many photon-photon impacts which tends to make the very thin layer of photons uniform eventually. Not all of these photon-photon[14] or gamma-gamma scattering's are elastic, many of these impacts emit radiation which reaches us as the cosmic microwave background the CMB.

Little ones merge to make big ones. In egg land, two eggs touch. Their shells merge much like soap bubbles merge to make a larger soap bubble. Their contents merge. Where there was two eggs, there is now one larger egg with two merged yokes.

Over time, there is a very big egg with many yokes merged together. The yokels, being unaware of the mechanics of merging, make up odd stories of their creation and their importance to the creator.

12 Large Scale Structure

There are groupings of mass in space so great, that gravity in the age of the Cosmos, would be inadequate for their formation from hydrogen gas. These are called large-scale structure [15]. An example is the Sloan great Wall [16]. Their great mass would have been reflected in the observed inhomogeneities in the CMB were the standard theory right.

The merging of black holes, does however, explain these structures and the Planck satellite asymmetry. The smaller black hole has a much higher density. The merged contents are enclosed in a much larger volume. The merged photons are bunched together. From within, one sees only the merged contents. The smaller black hole leaves behind a higher residual mass density, in the stretched out, merged contents, which is the artifact or footprint of their merging.

13 Spherical caps of merging spheres

See the figure above. A spherical cap is a part cut off a sphere. When two spheres merge they create a lens shaped merged region. The volume of the lens shaped merged region includes twice the volume of the spherical caps of each sphere. The volume of a spherical cap is,

$$\frac{1}{3} \pi * r^3 * (3 - fr) * fr^2 \quad (13.1)$$

with fr being the fraction of r , that is the height of the cap. The volume of a sphere equal to four spherical caps would be

$$\frac{4}{3}\pi * r^3 = \frac{4}{3}\pi * r^3 * (3 - fr) * fr^2 \quad (13.2)$$

$$1 = (3 - fr) * fr^2 \text{ therefore } fr = .6527036 \quad (13.3)$$

When the spherical caps of merging black holes of the same size reach .6527 of their radius, the volume and the mass of the merged portions satisfies the mass/radius formula for a black hole. The new velocity distribution in the merged black hole will cause all the orbits to relocate over time but these are small acceleration forces in a low density Cosmos like our own. Light and energy will eventually occupy a circular orbit at the new now larger radius of the black hole. The masses within will seek their own new orbits. The acceleration at the edge of the Cosmos is:

$$\frac{c}{age} = 6.33E-10 \frac{m}{s^2} \quad (13.4)$$

and is fractionally smaller within the Cosmos. This is billions of times smaller than the acceleration of gravity at the Earths surface of $9.8 \frac{m}{s^2}$. Far from being a dramatic event, the merging of low density black holes would be hardly detectable from an acceleration standpoint. A black hole approaching ours would be invisible until the merging. Its contents would then become visible in our Cosmos.

14 Comparing Traditional Black Holes and Orbiting Light Black Holes

There are two black hole formulas. First, in the orbiting light black holes we have rest energy equals gravitational energy and centrifugal force equals gravitational force. Second, in the traditional black holes we have kinetic energy equals gravitational

energy. The original black hole definition goes back centuries and is based on escape velocity from a star exceeding the speed of light. If the escape velocity only equals the speed of light then light will escape. This is the case with the Schwarzschild black hole. The idea with escape velocity is that the escaping object is slowing continuously and reaches infinity with zero velocity and zero kinetic and gravitational energy.

$$.5 * m * vr^2 = \frac{G * m * M}{r} \quad (14.1)$$

The kinetic energy equals the gravitational energy. Here vr is the radial velocity, the escape velocity. Substitute $vr = c$.

$$m * c^2 = 2 * \frac{G * m * M}{r} \quad (14.2)$$

The rest energy equals twice the gravitational energy. The energy of the escaping light is twice the gravitational energy.

$$\frac{m * c^2}{r} = 2 * \frac{G * m * M}{r^2} \quad (14.3)$$

Multiplied by $1/r$. The centrifugal force now equals twice the gravitational force. Light can not be restrained to an orbit.

$$r = \frac{2 * G * M}{vr^2} \quad (14.4)$$

Isolated r . If $r = \text{infinity}$ in this equation then

$$\text{infinity} = \frac{2 * G * M}{vr^2} \quad (14.5)$$

which is OK. vr is a variable which can approach zero at infinity.

$$\text{infinity} = \frac{2 * G * M}{c^2} = \text{Schwarzschild radius} \quad (14.6)$$

When $vr = c$ this is the Schwarzschild radius where it is frequently seen in General Relativity as $r=2M$ by using "geometric units" such that c and G are one [17]. Obviously, this odd

shorthand obscures the possible and necessary variability of G in an expanding and dynamic Cosmos. "Atomic units", where the mass, radius, charge, energy and Planck's constant within atoms are all one, also obscure variability.

If $r = \text{infinity}$ and $vr = c$ in (14.5) and (14.6) then

$$\text{infinity} = \frac{2 * G * M}{c^2} \quad (14.7)$$

which is false, c can not go to zero at infinity. Multiply by $\frac{.5 * m * c^2}{r}$.

$$.5 * m * c^2 = \frac{G * m * M}{r} \quad (14.8)$$

This is kinetic energy equals gravitational energy or the rest energy equals twice the gravitational energy. c is the escape velocity. Multiply by $\frac{2 * r}{m}$. These (14.9 - 14.11) are the traditional black hole equations.

$$c^2 * r = 2 * G * M \quad (14.9)$$

$$\frac{c^2 * r}{G * M} = 2 \quad (14.10)$$

$$r = \frac{2 * G * M}{c^2} = \text{Schwarzschild radius} \quad (14.11)$$

This radius is twice the value of the orbiting light black hole radius.

$$\frac{M}{r} = \frac{c^2}{2G} = 6.73E26 \frac{kg}{m} = \frac{\text{mass}}{\text{radius}} \text{ratio} \quad (14.12)$$

This is half the value of the orbiting light black hole.

$$M = \frac{c^2 * r}{2G} = 9.56E52 \text{ kg} = \text{mass of the Cosmos} \quad (14.13)$$

This is half the value of the orbiting light black hole.

$$\text{density} = \frac{\text{mass}}{\text{volume}} = \frac{3 * \text{mass}}{4\pi * r^3} = \frac{3c^2}{8\pi G r^2} \text{ so} \quad (14.14)$$

$$\text{density of the Cosmos} = \frac{3}{8\pi * G * r^2} = 7.97E-27 \frac{kg}{m^3} \quad (14.15)$$

This density of the Cosmos in traditional black holes decreases with age^2 like a surface, not with age^3 like a volume. Another hint that G must vary. Density is about 5 protons per meter cubed. This is half the density of the Cosmos we calculated for orbiting light black holes.

15 Tests for Orbiting Light Black Holes

Energy in orbit black holes are not the photo-sphere, within the event horizon of the Schwarzschild black hole.

The radius of traditional black holes is twice that of energy in orbit black holes. This difference may be detectable with the measurement of orbital periods of x-ray emitting clouds that orbit some black holes in binary systems of a black hole and star. A radiating mass of gas orbiting a black hole is like a lighthouse beacon sweeping past Earth hundreds of times per second. The orbital velocity of gas clouds at the Schwarzschild radius in traditional black hole is c which is not possible but in orbital light black holes the velocity at that radius is $.707*c$ and possibly detectable. The massive black hole at the center of our galaxy is expected to be impacted by an approaching gas cloud in the next year therefore we can look forward to the results.

16 Rotation of the Cosmos

As the light and energy orbit the expanding Cosmos, it takes longer to reach a reference point against the background universe. Newton would call this reference point absolute space. Mach would call it the fixed stars. The Cosmos, galaxy and solar system all rotate, with respect to that which is outside

Figure 4: Binary rotations



our cosmic dynamic unit. Looking at figure [4]. If the background universe has features which are close enough, and these features are not black holes, then they may be visible through the intense orbiting energy and light around the Cosmos. Seeing through this orbital energy seems possible since stars are visible near the sun during an eclipse as starlight perpendicular to the huge energy flow from the sun. We might be seeing such features in the Hubble telescope deep field photographs. It would not be remarkable if the background universe looks the same as it does within our dynamic unit Cosmos.

Angular Velocity

$$\frac{d}{dt} \text{angle of rotation} = \text{angular velocity} = \frac{\text{tangent velocity}}{\text{radius}} = \frac{vt}{r} = \frac{c}{c * \text{age}} = \frac{1}{\text{age}} = \frac{\text{radians}}{\text{second}} \quad (16.1)$$

$$\frac{d}{dt} \text{angle of rotation} = \frac{1}{\text{age}} \quad (16.2)$$

and

$$\frac{d}{dt} \ln(\text{age}) = \frac{1}{\text{age}} \quad (16.3)$$

so the

$$\text{angle of rotation} = \ln(\text{age}) = \text{natural logarithm of age} \quad (16.4)$$

$$\ln(4.73E17) = 40.7 \text{ radians} \quad (16.5)$$

The base of the natural logarithms is e.

$$e^{\text{angle of rotation}} = e^{40.7} = 4.73E17 = \text{age} \quad (16.6)$$

$$\ln(\text{age} * 2) = \ln(\text{age}) + \ln(2) \quad (16.7)$$

Each time the Cosmos doubles in age or size the angle of rotation of the Cosmos increases by

$$\ln(2) = .693 \text{ radians} = 39.7 \text{ degrees} \quad (16.8)$$

We are currently at 40.7 radians so

$$\frac{40.7}{2\pi} = 6.5 \text{ revolution} \quad (16.9)$$

might have been made by the orbiting light and energy in the age of the Cosmos. This revolution started when the Cosmos was only

$$e^{(40.7-6.28)} = 8.88E14 \text{ s} = 28.1 \text{ million years old} \quad (16.10)$$

We are currently at

$$e^{(40.7)} = 4.73E17 \text{ s} = \text{age} = 15 \text{ billion years old} \quad (16.11)$$

The next revolution will start in

$$e^{(40.7+6.28)} = 2.53E20 \text{ s} = 8017 \text{ billion years} \quad (16.12)$$

The slowly stirring Cosmos is slowing down.

$$\text{angular acceleration} = \frac{d}{dt} \frac{1}{\text{age}} = \frac{-1}{\text{age}^2} = -4.46E-36 \frac{1}{\text{s}^2} \quad (16.13)$$

This is the second derivative of the angle of rotation. This very small rate that the Cosmos is decelerating in its rotation is necessary for the equilibrium between rotation and expansion of the Cosmos. It satisfies the rules of dynamics. We are rotating with

the Cosmos. Everything has the same universal angular velocity $\frac{1}{age} = 2.11E-18\frac{1}{s}$ as a component of their local angular velocity as we will see in our galaxy. The Cosmos rotated faster when it was smaller and younger. This difference in rotation might be detected but the angular acceleration is profoundly slow at $\frac{-1}{age^2}$

Inertial accelerations

To calculate the path of expansion of a particle we need the vector sum of three accelerations; the centrifugal, tangent and coriolis. These are components of the so called fictitious forces which are more properly called forces due to inertia. They are certainly not fictitious if you take the Machian view [18] that inertia is the acceleration dependent gravitational force exerted by the rest of the Cosmos.

Centrifugal acceleration

The centrifugal and gravitational forces are equal. $\frac{m*c^2}{r}$ equals the centrifugal force and $\frac{c^2}{r}$ is the tiny acceleration felt by light or energy in orbit at the perimeter of the Cosmos.

$$\frac{c^2}{r} = \frac{c}{age} = 6.33\cdot 10\frac{m}{s^2} \quad (16.14)$$

Tangent Deceleration

We can calculate the tangent deceleration using the torque formula.

$$torque = moment\ of\ inertia * angular\ acceleration =$$

$$force * radius = m * a * r = .$$

$$m * r^2 * angularacceleration = m * a * r =$$

$$a = r * angularacceleration = tangent\ deceleration =$$

$$a = c * age * \frac{-1}{age^2} = \frac{-c}{age} \quad (16.15)$$

The direction of deceleration is opposite of rotation.

A note on the unknown moment of inertia

$$\text{moment of inertia of a ring} = mass * r^2 \quad (16.16)$$

$$\text{moment of inertia of a spherical shell} = \frac{2}{3} mass * r^2 \quad (16.17)$$

$$\text{moment of inertia of a sphere} = \frac{2}{5} mass * r^2 \quad (16.18)$$

We are not too far off when we call the Cosmic

$$\text{moment of inertia} = mass * r^2 \quad (16.19)$$

Forces perpendicular to the velocity

Centrifugal force = $\frac{m*vt^2}{r}$ is a radial force produced by a tangent velocity. The force is perpendicular to the velocity.

Deceleration force = $\frac{m*vr^2}{r}$ is a tangent force produced by a radial velocity. The force is perpendicular to the velocity.

Coriolis acceleration

Inertia will cause an outward directed mass, on a rotating platform, to lag behind in a direction opposite to the rotation. This is the reaction. The action which is the coriolis acceleration is in the direction of the rotation. A person in an accelerating car is pushed back against the seat. This is a reaction to the acceleration. The acceleration is in the direction of the velocity. The reaction is in the direction opposite the velocity.

$$\text{coriolis acceleration} = 2 * \text{angular velocity} * vr =$$

$$2 * \frac{vt}{r} * vr = 2 * \frac{c}{c * age} * c = 2 * \frac{c}{age} \quad (16.20)$$

vr, the radial velocity at the perimeter of the Cosmos, is c.

17 Cosmic Expansion

Now that we have calculated the inertial accelerations, we can look at the way the Cosmos expands. We have the

$$\text{centrifugal acceleration} = \frac{c}{age} \quad (17.1)$$

directed radially out. We have the

$$\text{coriolis acceleration} = 2 * \frac{c}{age} \quad (17.2)$$

in the direction of rotation, and the

$$\text{rotational deceleration} = \frac{-c}{age} \quad (17.3)$$

in the direction opposite of rotation. The resultant of these accelerations, is 45 degrees between the direction of rotation and the outward directed radius. It has a value of,

$$\sqrt{2} * \frac{c}{age} = 8.96E-10 \frac{m}{s^2} \quad (17.4)$$

A particle moving in this way traces out a logarithmic spiral.

We have seen that the

$$\text{angle of rotation} = \ln(age) \quad (17.5)$$


This can be written as

$$age = e^{(\text{angle of rotation})} \quad (17.6)$$

Now

$$r = c * age \quad (17.7)$$

Figure 5: Logarithmic spirals



can be written as

$$r = c * e^{(angle\ of\ rotation)} \quad (17.8)$$

This is the equation of a logarithmic spiral. It is no coincidence that many galaxies have a spiral shape. Their spiral is expected in an expanding rotating Cosmos.

Logarithmic Spiral

Looking at figure [5]. Indeed, it is not that space expands, but that the distance between orbiting masses increases as they spiral out and apart from each other as the Cosmos expands and slows in its rotation. The tangent velocity of the stars orbiting in galaxies, stays the same as the galaxies expand and the orbital periods increase per the flat rotation curves of galaxies.

Any velocity change would require force and energy which are absent.

Torque

*torque = moment of inertia * angular acceleration =*

$$M * r^2 * \text{angular acceleration} =$$

$$M * vr^2 * age^2 * \text{angular acceleration}$$

but

$$\frac{vr}{c} = \frac{M}{Mc} \quad \text{and} \quad vr^2 = \frac{c^2 * M^2}{Mc^2}$$

therefore

$$M * \frac{c^2 * M^2}{Mc^2} * age^2 * \text{angular acceleration} =$$

$$\frac{c^2 * M^3}{Mc^2} * age^2 * \text{angular acceleration} = \text{torque}$$

If the mass of the black hole is $M = Mc$ the mass of the Cosmos then

$$\text{torque} = Mc * c^2 * age^2 * \frac{-1}{age^2} = Mc * c^2 \quad (17.9)$$

We see that the age^2 in the square of the radius, in the moment of inertia, increases at the same rate the angular acceleration $1/age^2$ decreases so that the age^2 in each cancels and the energy stays constant. We will see the same thing in the torque of a spinning galaxy. The radius of the Cosmos increases while the rotation of the Cosmos slows down, and with it all the black holes and galaxies, without a change in energy or use of power, always in dynamic equilibrium. Orbits spiral out as the gravitational force decreases with the age of the Cosmos.

18 Galactic Matters

Dark matter can be explained by matter present farther out from the galaxy than is seen in the optical or radio spectrum. There

are orbiting atomic hydrogen clouds in this dark region. Each meter of radius adds a fixed amount of mass to a bigger volume so the density decrease with radius. The galaxy extends to the radius at which the galaxy is the same density as the Cosmos. We have a tangent velocity vt , of 210,000 m/s [19] at a radius within the galaxy of 30 kpc = 98,000 light years = 9.26E20 m. The mass contained within this orbit is the visible matter

$$visible\ matter = mass = \frac{vt^2 * radius}{G} =$$

$$6.12E41\ kg\ or\ 308\ billion\ solar\ masses \quad (18.1)$$

The mass/radius ratio

$$\frac{mass}{radius} ratio = \frac{m}{r} = \frac{vt^2}{G} = \frac{6.12E41\ kg}{9.26e20\ m} = 6.61E20 \frac{kg}{m} \quad (18.2)$$

The mass added by the next meter of radius, 6.61E20 kg, when divided by the volume added by the next meter of radius, is the density at this radius, or the mass/radius ratio,

$$\frac{m}{r} = \frac{vt^2}{G} \quad (18.3)$$

divided by the surface area of a sphere at this radius.

$$\frac{6.61E20 \frac{kg}{m}}{4\pi * 9.26E20^2 m^2} = 6.13E-23 \frac{kg}{m^3} \quad (18.4)$$

I suspect that this low density of matter or dark matter, would usually be hard to detect, for example with 21 cm radiation[20], but it is obviously still probably matter not some mysterious stuff. Radio telescopes can detect the atomic hydrogen at 21 cm, if it is dense enough along their line of sight. Cold molecular hydrogen [21] which is more stable and probably much more common is unfortunately invisible at radio wavelengths. It may be detected in the future as the unseen dark matter.

19 Flat Rotation Curves of Galaxies

$$\frac{m * vt^2}{r} = \text{frac}G * m * Mr^2 \quad (19.1)$$

Using (2.1) in reference to galaxies. Multiply by r^2/m .

$$vt^2 * r = G * M \quad (19.2)$$

As r the radius within a galaxy increases, vt^2 the tangent velocity of stars, at that radius within the galaxy should decrease with $G*M$ taken as a constant, in keeping with Kepler. However, what is seen is that the tangent velocity vt is largely flat, that is, the velocity stays the same with increasing radius once outside the galactic core. This is called a flat rotation curve[22] as illuminated by Vera Rubin[23]. These common flat rotation curves have been used as evidence that there is dark matter in galaxies. There is another more sensible conclusion. The above equation can be written as:

$$\frac{M}{r} = \frac{vt^2}{G} = \frac{\text{mass}}{\text{radius}} \text{ratio} \quad (19.3)$$

The vt will stay constant with increasing radius if the mass/radius ratio is maintained. This is similar to the mass/radius ratio seen as a defining condition for a black hole and the Cosmos.

$$\frac{M}{r} = \frac{c^2}{G} = 1.35E27 \frac{\text{kg}}{m} = \frac{\text{mass}}{\text{radius}} \text{ratio} \quad \text{from (2.6)}$$

Galactic Density

The galaxy density increases with a constant m/r where

$$\frac{m}{r} * R = \text{mass} \quad \text{and} \quad \frac{3}{4\pi * R^3} = \text{volume} \quad (19.4)$$

$$\frac{m * 3 * R}{r * 4\pi * R^3} = \frac{m * 3}{r * 4\pi * R^2} = \text{galactic density} \quad (19.5)$$

Figure 6: Galaxy expansion



The galactic density decreases as $1/R^2$. I postulate that the upper limit for the radius of the galaxy would be the radius at which the galactic density at that radius R equals the average density of the Cosmos

$$\frac{m * 3}{r * 4\pi * R^2} = \frac{3 * Mc}{4\pi * c^3 * age^3} = \text{Cosmic density} \quad (19.6)$$

but $m/r = vt^2/G$ so

$$\begin{aligned} \frac{vt^2}{G * R^2} &= \frac{Mc}{c^3 * age^3} \\ \frac{vt^2}{R^2} &= \frac{Mc * G}{c^3 * age^3} \\ \frac{vt^2}{R^2} &= \frac{c^2 * c * age}{c^3 * age^3} \\ vt^2 * age^2 &= R^2 \\ vt * age &= R \end{aligned} \quad (19.7)$$

20 Galactic Expansion

Galactic expansion is seen in figure [6]. The tangent velocity vt , which is seen in the flat rotation curves of galaxies, times the age of the Cosmos equals the radius of the galaxy. This suggest that there is a Hubble expansion occurring within the galaxy.

$$R = vt * age = 210,000 \frac{m}{s} * 4.73E17 s = 9.94E22 m = 3.22 Mpc$$

$$R = 10.5 \text{ million light years} = \text{galactic radius} \quad (20.1)$$

The galaxy is as low in density as the Cosmos at the perimeter of the galaxy. Hubble's constant = $H_0 = 65 \text{ km}/(\text{s} * \text{Mpc})$

$$R * H_0 = 3.22 \text{ Mpc} * 65 \frac{\text{km}}{\text{s} * \text{Mpc}} = 209,000 \frac{\text{m}}{\text{s}} = vr \approx vt \quad (20.2)$$

The radial velocity equals the tangent velocity. When the tangent velocity of something equals its radial velocity. It spirals out at a constant angle of 45 degrees. This is the same spiral as the Cosmos.

21 Dark matter

$$Mg = \text{galactic mass} = \frac{m}{r} * R$$

$$Mg = 6.61E20 \frac{\text{kg}}{\text{m}} * 9.94E22 \text{ m} = 6.57E43 \text{ kg} \quad (21.1)$$

Dark matter = total matter - visible matter within 30 kpc which is $6.51E43 \text{ kg}$

$$\text{Dark matter} = 6.57E43 \text{ kg} - 6.12E41 \text{ kg} = 6.51E43 \text{ kg} \quad (21.2)$$

$$\frac{6.51E43 \text{ kg}}{6.57E43 \text{ kg}} = \frac{\text{dark matter}}{\text{visible matter}} \text{ratio} = \frac{106}{1} \quad (21.3)$$

The 106/1 ratio of dark matter/visible matter.

$$\frac{vt^2}{G} = \frac{Mg}{r} \quad (21.4)$$

For vt^2 and M/r to remain constant,

$$\frac{vt^2 * Mc}{c^3 * \text{age}} = \frac{Mg}{vt * \text{age}} \quad (21.5)$$

substituted $G = c^3 * \text{age}/Mc$, $r = vt * \text{age}$ and $Mc = c^3/gk$.

$$\frac{vt^3}{c^3} = \frac{Mg}{Mc} \text{ or } \frac{vt^3}{c^3} = \frac{Mg * gk}{c^3} \quad (21.6)$$

The ratio of velocities and masses.

$$Mg = \text{mass of galaxy} = \frac{Mc * vt^3}{c^3} = \frac{vt^3}{gk} \quad (21.7)$$

Compare this with (4.7) where the mass of the Cosmos = c^3/gk [7]

Torque of the spinning galaxy

Here mass is the mass of the galaxy. r is the radius of the galaxy. vr is the radial velocity of expansion at the perimeter of the galaxy or $vr = vt$ the characteristic tangent velocity of the flat rotation curve of the galaxy.

*torque = moment of inertia * angular acceleration =*


$$\begin{aligned} & \text{mass} * r^2 * \text{angular acceleration} = \\ & \text{mass} * vr^2 * age^2 * \frac{1}{age^2} = \text{mass} * vr^2 \end{aligned} \quad (21.8)$$

We see that the square of the radius in the moment of inertia for the galaxy, $vr^2 * age^2$ increases at the same rate the angular acceleration of the galaxy $1/age^2$ decreases so that the age^2 in each cancels and the energy stays constant. The radius of the galaxy increases while the rotation of the galaxy slows down without a change in energy or use of power. Orbits spiral out as the gravitational force decreases with the age of the Cosmos.

22 Hubble expansion in the Solar System

If the Hubble expansion extends to the solar system as shown in figure [7], then all the planets share the same very small precession[24] rate, of their major axis within the planet's elliptic plane. An expanding ring slows in its rotation. We calculate a

Figure 7: Solar system expansion



change in angular velocity which is due to a radial Hubble velocity. This effect may be detected by atomic clocks or by large ring laser gyroscopes [25] which detect absolute rotations.

The length of the orbit, when divided by 2π radians per revolution equals the meters per radian or

$$\text{meters per radian} = \frac{2\pi * r}{2\pi * \text{radians}} = \frac{r}{\text{radians}} \quad (22.1)$$

r is the distance from the sun to any planet. $vr = r/\text{age}$, the radial velocity, the Hubble expansion velocity in meters per second.

$$\frac{vr}{\text{meters per radian}} = \frac{\frac{r}{\text{age}}}{\frac{r}{\text{radians}}} = \frac{\text{radians}}{\text{age}} \quad (22.2)$$

We see that the r 's cancel so that this rate of precession is universally true for the entire solar system and is not tied to any radius. This is Hubble as a rotation rate.

$$\begin{aligned} \frac{\text{radians}}{15 \text{ billion years}} &= 6.66E^{-11} \frac{\text{radians}}{\text{year}} = \\ 1.375E^{-5} \frac{\text{arcs}}{\text{year}} &= \text{precession} \end{aligned} \quad (22.3)$$

This is a very small angle to measure. The gravity probe B satellite was seeking to measure a Lense-Thirring frame dragging of $4,200E^{-5}$ arcs/year.

$$\frac{31,556,926 \frac{\text{seconds}}{\text{year}}}{1,296,000 \frac{\text{arcs}}{\text{year}}} = 24.35 \frac{\text{seconds}}{\text{arc}} \quad (22.4)$$

for the Earth to cover one arc second of circumference in its orbit.

$$24.35 \frac{\text{seconds}}{\text{arc}} * 1.375E^{-5} \frac{\text{arcs}}{\text{year}} = 3.35E^{-5} \frac{\text{seconds}}{\text{year}} \quad (22.5)$$

which is the amount added yearly to the orbital period by the Hubble expansion. This is one leap second being added to our orbit in the solar system every 2,986.8 years. The entire solar system is slowing in its rotation while it expands, like a dynamic unit, like the galaxy and like the Cosmos. It is a consequence of the slowing rotation of the Cosmos and is tied by dynamics to the expansion of the Cosmos. The precession is proportional to $1/\text{age}$.

$$vr = \text{radial velocity} = \frac{r}{\text{age}} \quad (22.6)$$

The Earth orbits the sun at 149E9 m and

$$vr = \frac{149E9 \text{ m}}{4.73E17 \text{ s}} = 3.16E^{-7} \frac{\text{m}}{\text{s}} = 9.97 \frac{\text{m}}{\text{year}} \quad (22.7)$$

vr is too small to measure for the Earth. The moon orbits the Earth much closer at 380E6 m but the Apollo missions to the moon left behind laser reflectors so that the round trip of laser pulses could be timed and the receding velocity of the moon measured.

$$vr = \frac{380E6 \text{ m}}{4.73E17 \text{ s}} = 8.02E^{-10} \frac{\text{m}}{\text{s}} = 25.3 \frac{\text{mm}}{\text{year}} \quad (22.8)$$

The moons velocity is 38 mm/year and is usually attributed to tidal drag and not the expansion of the universe. I postulate, 25 mm is due to a Hubble expansion and 13 mm is due to tidal drag.

Planetary Spirals

To calculate the path of expansion of a planet we need the vector sum of three accelerations. We have the

$$\text{centrifugal acceleration} = \frac{vr}{age} \quad (22.9)$$

directed radially out. We have the

$$\text{coriolis acceleration} = 2 * \frac{vr}{age} \quad (22.10)$$

in the direction of rotation, and the

$$\text{tangent rotational deceleration} = \frac{-vr}{age} \quad (22.11)$$

in the direction opposite of rotation. The resultant of these accelerations, is 45 degrees between the direction of rotation and the outward directed radius. It has a value of,

$$\sqrt{2} * \frac{vr}{age} \quad (22.12)$$

A particle moving in this way traces out a logarithmic spiral.

$$\text{torque} = \text{moment of inertia} * \text{angular acceleration} =$$

$$m * r^2 * \text{angular acceleration} =$$

$$m * vr^2 * age^2 * \text{angular acceleration} =$$

$$m * vr^2 * age^2 * \frac{-1}{age^2} = \text{mass} * vr^2 \quad (22.13)$$

Here mass is the mass of a planet. r is the distance from the sun. $vr = r/age$ is the radial velocity. We see that the square of the radius in the moment of inertia for the planet, $vr^2 * age^2$ increases at the same rate the angular acceleration of the planet, $1/age^2$ decreases so that the age^2 in each cancels and the energy stays constant. Expansion and rotation rates are linked. The distance from the sun to the planets increases and the orbital periods of the planets lengthen without a change in velocity, energy or the use of power. Orbits spiral out as the gravitational force decreases with the age of the Cosmos.

23 Ellipses

The ellipse is the path of point that moves, so that the sum of its distance from the two foci is constant. A whisper at one focus of an elliptical room, is heard at the other focus in the room, because the distance and travel time from focus to focus, is the same for any path of sound reflecting off the walls. Light acts the same, with the same geometry, with ellipsoidal mirrors. The two foci can be called the origin and the destination. Light leaves the origin as an expanding sphere and reflects at the ellipsoidal surface as a ring. It is a ring because it is the intersection of a sphere and ellipsoid. This reflection focuses the light on to the destination. The overall travel time from the origin to reflection, to destination, is always the same for all angles of light departing from the origin. If we retain the origin and destination and change to a luminous expanding sphere we can eliminate the mirror and reflection while keeping the geometry of the intersection of an expanding sphere and ellipsoid. The luminous sphere is the shell of Cosmos which emits the CMB. Now the ellipsoid is an abstraction that exists in concept but not in reality. The ellipsoid defines the path of things that are perceptible at our location at a certain time.

24 Cosmic Cross Section

$$r = \frac{c * age}{2} * \frac{1 - fr^2}{1 + \cos(angle) * fr} \quad (24.1)$$

A cosmic cross section is seen in figure [8]. We are located at someplace like d, at an unknown fraction of the radius of the Cosmos = fr, on a concentric radius = fr*c*age and are moving radially and tangentially with a velocity = fr*c. This is the ellipse drawn from the center focus, which is the origin, o on figure [8]. The radius of the expanding sphere, which intersects the ellipse is r in the equation. As the angle is varied, the points form an ellipse stretched in the radial direction with fr*c*age as

Figure 8: Cosmic cross section




Figure 9: Expanding ellipses



the distance between the foci, od on figure [8]. The sum of the distances from the two foci, to a point, is always equal to c^* age. The angle and radius when rotated around the center line trace a ring that is the intersection of the sphere and the ellipsoid, aa or bb or cc on figure [8]. This nested series of rings may partially polarize the CMB, along the axis of the origin. Other polarizations seem likely. There is evidence that light traveling through space is polarized in a non-random direction [26]. There is also evidence from the Planck satellite [27] that space could be rotating from the asymmetry in the average temperatures of the CMB on opposite hemispheres of the sky as if we are offset from the center and seeing differences in velocity as differences in temperature.

25 Blackbody watts from temperature

A flux of radiation has a Kelvin temperature. We see the temperature of the sun as 5778 Kelvin. We can convert this to $\frac{\text{watts}}{\text{meter}^2}$ for black body radiation with the Stephan and Boltzmann law.

$$K^4 = 5.5698E^{-8} \frac{W}{m^2 * K^4} = \frac{W}{m^2} \quad (25.1)$$

$$\text{The suns temperature} = 5778 \text{ Kelvin} = 62E6 \frac{W}{m^2} \quad (25.2)$$

The sun has a wattage of $62E6 \frac{\text{watts}}{\text{meter}^2}$ over the surface area of the sun of $4\pi * (695E6 \text{ m})^2$ or a total of 3.77E26 watts. At the

distance of the Earth, 149E9 m, this is

$$\frac{3.77E26 \text{ watts}}{4\pi * (149E9 \text{ m})^2} = 1341.8 \frac{\text{watts}}{\text{m}^2} = 394 \text{ K} = 249.5 \text{ F} \quad (25.3)$$

Space is quite hot in the sun. Only the Earth's atmosphere protects us. Threat her with respect. We can work the CMB in just this way.

26 Uniformity of the CMB

The question of the uniformity of the CMB, is usually answered; that we appear to be at the center of the Big Bang, because the Big Bang explosion happened everywhere. A more quantitative, and less miraculous solution, looks again at figure [8]. A ray of CMB reaches us after two distinct intervals. The first interval starts at the center focus, at the origin, at point o, on figure [8]. It is with the expanding and cooling spherical shell, before the ray of CMB, which we will be observing, is emitted. Examples on figure [8] are the lines oa, ob and oc. The temperature and $\text{watts}/\text{meter}^2$, of the expanding sphere, is proportional to the inverse square of the radius. The second interval, is the travel of the ray of CMB through space, after it leaves the expanding spherical shell. Examples on figure [8]; are the lines ad, bd and cd. It ends with the reception of the CMB, at the observer at point d. The temperature and $\text{watts}/\text{meters}^2$, of the CMB during the second interval, is also proportional to the inverse square of the radius. The two intervals always add up to $c * \text{age}$ meters in age seconds in any direction the observer looks. For CMB emitted early in the Cosmos, there is a shorter interval with the expanding sphere, and a much longer path through space interval to reach the observer. For CMB emitted later, there is a longer spherical expansion interval, before the CMB is emitted, but a shorter path through space interval to the observer. The expanding spherical shell, the CMB which it emits, and the CMB during its travel through space, all have a temperature

proportional to the inverse square of the distance traveled from the origin o. When the radius of the Cosmos was oa, it emitted CMB from the entire spherical surface. Only that from the ring, on the sphere at aa, will reach d at the same time as the other rings on the same ellipsoid. A similar argument can be seen in the rings bb and cc. All the CMB from the various rings which intersect the ellipse arrive at point d at the same time and temperature.

$$oa + ad = ob + bd = oc + cd = or = c * age \quad (26.1)$$

The formulas in the next section, show the relationship between radius and temperature. The CMB had a temperature at point a of 5.4 K when the Cosmos was one forth its age and size. It arrived at point d at 2.7 K after expanding for three fourths the age of the Cosmos. The temperature at point b was 3.8 K when the Cosmos was one half its age and size. It arrived at point d at 2.7 K after expanding for half the age of the Cosmos. The temperature at point c was 3.2 K when the Cosmos was three fourths its age and size. It arrived at point d at 2.7 K after expanding for one forth the age of the Cosmos. The temperature is going down because of the inverse square law of radiation not because of the expansion of the vacuum of space.

27 Black body watts from the CMB

A flux of radiation has a Kelvin temperature. We see the temperature of the CMB as 2.735 Kelvin. We can convert this to *watts/meter*² for black body radiation with the Stephan and Boltzmann law.

$$K^4 = 5.5698E-8 \frac{W}{m^2} \quad (27.1)$$

$$CMB \text{ temperature} = 2.735 \text{ Kelvin} = 3.11E-6 \frac{W}{m^2} \quad (27.2)$$

When one sees something, it is in terms of W/m^2 and the inverse square law. The W/m^2 times the area of the Cosmos

= wattage of the CMB, because as we saw in figure [8], the temperature at point d, where the observer is located, is the same as a point r, the radius of the Cosmos.

$$3.11E-6 \frac{W}{m^2} * 4\pi * rc^2 = 7.9E47 W = power = \frac{energy}{second} \quad (27.3)$$

The CMB has the luminosity of a 7.9E47 watt light bulb seen from a distance of 15 billion light years. This is the same as 7.9E47 watts stretched over the area of a sphere with a radius of 15 billion light years. The CMB is emitted from the expanding radiant shell which is where light orbits at the perimeter of the Cosmos. The non-elastic impact of the photons within this thin spherical shell is the source of the CMB. The light was accumulated in this shell as the Cosmos gained mass and light through the merging of black holes. The energy emitted in 15 billion years by the CMB, if the energy output is constant, is

$$7.9E47 \text{ watts} * age = 3.7E65 \text{ Joules} \quad (27.4)$$

For comparison, the energy of the Cosmos,

$$Mc * c^2 = 1.7E70 \text{ Joules} \quad (27.5)$$

$$\frac{Mc * c^2}{3.11E-6 \frac{W}{m^2} * 4\pi * c^2 * age^2} = \frac{1.7E70 J}{3.7E65 J} = 46,000 \quad (27.6)$$

If the CMB is the remnant energy from the Big Bang then why is it so feeble? However, if the CMB is emitted instead through the non-elastic impact of photons at the perimeter of the Cosmos then this small value of energy makes some sense. The gravitational and centrifugal accelerations on the photon in orbit are c/age . As the Cosmos expands the photons orbit at a larger radius and the orbital accelerations decrease. The rate of change of the acceleration is $1/age^2$. The W/m^2 of the CMB is:

$$CMB = \frac{7.9E47 W}{4\pi * rc^2} = \frac{7.9E47 W}{4\pi * c^2 * age^2} = \frac{W}{m^2} \quad (27.7)$$

The rate of change of the orbital acceleration is proportional to the W/m^2 of the CMB. We can map the power of the CMB onto the smaller spheres and higher temperature when the Cosmos was younger, as long as we keep well clear of infinities.

$$\frac{\frac{7.9E47 W}{\sqrt{4\pi*5.569E-8}}}{(\text{temperature } K)^2} = \text{radius } m \quad (27.8)$$

Collect terms.

$$\frac{1.0615E27 m}{(\text{temperature } K)^2} = \text{radius } m \quad (27.9)$$

The radius, of the expanding sphere of the shell, is proportional to the inverse square of the temperature. The following examples map temperature and radius.

The radius of the Cosmos currently at 2.735 K.

$$\frac{1.0615E27 m}{2.735 K^2} = 1.42E26 m =$$

$$1.42E26 m = c * \text{age} = rc = 15 \text{ billion light years} \quad (27.10)$$

At an age of 1.5 million years, at the freezing point of water of 273 K.

$$\frac{1.0615E27 m}{273 K^2} = 1.42E22 m =$$

$$1.42E22 m = \frac{rc}{10000} = 1.5 \text{ million light years} \quad (27.11)$$

At a temperature below 3000 K plasma becomes transparent to light. The radius becomes

$$\frac{1.0615E27 m}{3000 K^2} = 1.18E20 m =$$

$$1.18E20 m = 12,466 \text{ light years} \quad (27.12)$$

At an age of 12466 years.

28 What is reasonable?

This last examples would fit in the core of a galaxy. This is only the CMB, but this much power would require an absurdly large star. An inside out or hollow star since its radiation comes to us from every direction. We have allowed the ease of doing calculations to project something absurd. We can see that these formulas and others like them might be used to trace back to a creation event at a point of infinite temperature and density. This has become dogma, trussed up with patches, which helps obscure the absurdity of physical infinities. At its present mass, our Cosmos could never have been that small or young. Since little ones make big ones, the Cosmos came about by the merging of black holes. Low density black holes are big, old and expanding fast so they incorporate a lot of space over time. Big ones present a bigger target for merging. Old ones present a target for merging that has been around for a long time. Space seems well populated with black holes. It is a small step, for our dynamic unit, our Cosmos, to be just another ordinary low density black hole in a universe full of the same. A ledger might have beliefs on the left side, and evidence for those beliefs on right side. The dynamics described here are mathematically consistent beliefs, which don't require physical infinities. The evidence is the values presented by the mass, radius and density of the Cosmos, source and uniformity of the CMB, the spiral shape and the flat rotation curves of galaxies and the prevalence of dark matter. All the parts slip together seamlessly, and the dynamics locks all the parts together. There are no free parameters which might be adjusted to reflect a point of view.

References

- [1] Edwin Hubble @ http://en.wikipedia.org/wiki/Hubble's_law
- [2] age of the Cosmos @ <http://hypertextbook.com/facts/1999/RachelHoover.shtml>
- [3] radius of the Cosmos @ <http://hypertextbook.com/facts/2002/CarmenBissessar.shtml>
- [4] non-Doppler redshift @ <http://newtonphysics.on.ca/hubble/index.html>
- [5] Geons @ J. A. Wheeler, Geons, Phys. Rev. 97:511-36 (1955)
- [6] mass without mass @ Ciufolini and Wheeler, Gravitation and Inertia, 77-78 (1995)
- [7] mass of the Cosmos @ <http://hypertextbook.com/facts/2006/KristineMcPherson.shtml>
- [8] Ciufolini and Wheeler @ Ciufolini and Wheeler, Gravitation and Inertia, 5 (1995)
- [9] expanding Cosmos @ <http://blackholeformulas.com/files/dconccirc.10.gif>
- [10] triangular definitions @ <http://blackholeformulas.com/files/dctri.10.gif>
- [11] Electric gravity @ <http://blackholeformulas.com/files/electricgravity.html>
- [12] kinetic energy @ <http://www.newtonphysics.on.ca/kinetic/index.html>
- [13] density of the universe @ <http://hypertextbook.com/facts/2000/ChristinaCheng.shtml>

- [14] @ http://en.wikipedia.org/wiki/Nuclear_density
- [15] neutron star mass @ http://www.lsw.uni-heidelberg.de/users/mcamenzi/NS_Mass.html
- [16] x-ray burster radius @ <http://www.astro.washington.edu/users/ben/a510/NSTARS.new.html>
- [17] photon-photon @ <http://www.hep.ucl.ac.uk/opal/gammagamma/gg-tutorial.html>
- [18] large scale structure @ http://en.wikipedia.org/wiki/Large-scale_structure_of_the_Cosmos
- [19] Sloan great wall @ http://en.wikipedia.org/wiki/Sloan_Great_Wall
- [20] $r=2M$ @ Jean-Pierre Luminet, Black Holes, 120 (1992)
- [21] Machian view @ <http://www.bun.kyoto-u.ac.jp/~suchii/mach.pr.html>
- [22] 210,000 m/s @ <http://www.astronomynotes.com/ismnotes/s7.htm>
- [23] 21 cm radiation @ http://en.wikipedia.org/wiki/Hydrogen_line
- [24] molecular hydrogen @ <http://www.newtonphysics.on.ca/hydrogen/index.html>
- [25] flat rotation curve @ http://www.astro.cornell.edu/academics/courses/astro201/rotation_curves.htm
- [26] Vera Rubin @ <https://en.wikipedia.org/wiki/VeraRubin>
- [27] precession @ <http://en.wikipedia.org/wiki/Precession>

[28] large ring laser gyroscopes @ <http://www.wettzell.ifag.de/LKREISEL/G/LaserGyros.html>

[29] polarized in a non-random direction @ <http://www.spie.org/web/oer/june/jun97/axis.html>

[30] Planck satellite @ http://www.esa.int/Our_Activities/Space_Science/Planck/Planck_reveals_an_almost_perfect_Universe

Go to Index @ <http://blackholeformulas.com>